
No.



200000215

THE UNITED STRATES OF AMIERIOS

Rijk Zwann Zandteelt en Zandhandel B.H.

MICTORS, THERE HAS BEEN PRESENTED TO THE

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY TUSIS UNOM THE DATE OF THIS GRANT SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC CILLERISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE WILLIAM EXCLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR TINGIT, OR EXPORTING IT, OR CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE TRPOSE, OR CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE ABOVE R USING IT IN PRODUCING A HYBRID OR DIFFERENT VARIETY THEREFROM, TO THE EXTENT THE PLANT VARIETY PROTECTION ACT. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

LETTUCE

'Toronto'

In Jestimonn Mercos, I have hereunto set my hand and caused the seal of the Hunt Hurichn Frotection Office to be affixed at the City of Washington, D.C. this seventh day of August, in the year two thousand and six.

Plant Variety Protection Offic

SCIENCE DIVISION - PLANT VARIETY PROTECTION OFFICE Application is required in order to determine if a plant variety protection certificate is to be issued (7 U.S.C. 2421) Information is held confidential until certificate is issued (7 U.S.C. 2426). APPLICATION FOR PLANT VARIETY PROTECTION CERTICATE (Instructions and information collection burden statement on reverse) 3. VARIETY NAME 1. NAME OF APPLICANT(S) (as it is to appear on the Certificate) 2. TEMPORARY DESIGNATION OR EXPERIMENTAL NUMBER TORONTO Rijk Zwaan Zaadteelt en Zaadhandel B.V. 4. ADDRESS (Street and No., or R.F.D. No., City, State and ZIP Code, and Country) 5. TELEPHONE (include area code) FOR OFFICIAL USE ONLY 70°11111110215 + 31 174 532300 Burgemeester Crezeelaan 40 2678 KX DE LIER, The Netherlands 6. FAX linclude area codel + 31 174 513730 FILING AND 7. GENUS AND SPECIES NAME 8. FAMILY NAME (Botenical) **EXAMINATION FEE** Lactuca sativa L. Compositae \$ J450 : 9. CROP KIND NAME (Common name) May 1,2000 Lettuce CERTIFICATION FEE: 10. IF THE APPLICANT NAMED IS NOT A "PERSON", GIVE FORM OF ORGANIZATION (corporation, partnership, association, etc. Rijk Zwaan Zaadteelt en Zaadhandel B.V. Corporation 11. IF INCORPORATED, GIVE STATE OF INCORPORATION 12. DATE OF INCORPORATION The Netherlands 08-18-1953 Corporation 13, NAME AND ADDRESS OF APPLICANT REPRESENTATIVES, IF ANY, TO SERVE IN THIS APPLICATION AND 14. TELEPHONE (include are RECEIVE ALL PAPERS codel 15. FAX (include area code) 16. CHECK APPROPRIATE BOX FOR EACH ATTACHMENT SUBMITTED (Follow instructions on reverse) a. Exhibit A. Origin and Breeding History of the Variety b. Exhibit B. Statement of Distinctness c. Exhibit C. Objective Description of the Variety d. Exhibit D. Additional Description of the Variety e. Exhibit E. Statement of the Basis of the Applicant's Ownership f. 🗷 Voucher Sample (2.500 viable untreated seeds or, for tuber propagated varieties verification that tissue culture will be deposited and maintained in a public repository) g. Filing and Examination Fee (\$2,450) made payable to the "Treasurer of the United States" (Mail to PVPO) 17. DOES THE APPLICANT SPECIFY THAT SEED OF THIS VARIETY BE SOLD BY VARIETY NAME ONLY, AS A CLASS OF CERTIFIED SEED (See Section 83(a) of the Plant Variety Protection Act!? ☐ YES (If "yes", answer items 18 and 19 below) ■ NO (If "no", go to item 20) 18. DOES THE APPLICANT SPECIFY THAT SEED OF THIS VARIETY BE LIMITED AS 19. IF "YES" TO ITEM 18, WHICH CLASSES OF PRODUCTION TO NUMBER OF GENERATIONS? BEYOND BREEDER SEED? □ NO ☐ FOUNDATION ☐ REGISTERED ☐ CERTIFIED ☐ YEŞ 20. HAS THE VARIETY OR A HYBRID PRODUCED FROM THE VARIETY BEEN RELEASED, USED, OFFERED FOR SALE, OR MARKETED IN THE U.S. OR OTHER COUNTRIES? ■ YES (if "yes", give names of countries and dates) □ NO Toronto has been sold for the first time in Australia in October 1999 and has not yet been sold in the U.S.A. 21. The applicant(s) declare that a viable sample of basic seed of the variety will be furnished with application and will be replenished upon request in accordance with such regulations as may be applicable, or for a tuber propagated variety a tissue culture will be deposited in a public repository and maintained for the duration of the certificate. The undersigned applicant(s) is(are) the owner(s) of this sexcually reproduced or tuber propagated plant variety, and believe(s) that the variety is new, distinct, uniform, and stable as required in Section 41, and is entitled to protection under the provisions of Section 42 of the Plant Variety Protection Act. Applicant(s) is(are) informed that false representati an jeopordize protection and result in penalties SIGNATURE OF APPLICANT (Owner(s)) SIGNATURE OF APPLICANT (Owner(s)) NAME (Please print or type) NAME (Please print or type) M. Zwaan CAPACITY OR TITLE DATE CAPACITY OR TITLE DATE Managing Director 18 April 2000

1974 (5 U.Š.C. 552a).

U.S. DEFARTMENT OF AGRICULTURE

AGRICULTURAL MARKETING SERVICE

GENERAL: To be effectively filed with the Plant Variety Protection Office (PVPO), ALL of the following items must be received in the PVPO: (1) Completed application form signed by the owner; (2) completed Exhibits A, B, C, E; (3) at least 2,500 viable untreated seeds, or for tuber reproduced varieties verification that a viable (in the sense that it will reproduce an entire plant) tissue culture will be deposited and maintained in a public repository prior to issuance of a certificate; (4) check drawn on a U.S. bank for \$ 2,450 (\$ 300 filing and \$2,150 examination fee), payable to "Treasurer of the United States" (See Section 97.175 of the Regulations and Rules of Practice.) partial applications will be held in the PVPO for not more than 30 days, then returned to the applicant as unfilled. Mail application and other requirements to Plant Variety Protection Office, AMS, UDA, Room 500, NAL Building, 10301 Baltimore Blvd., Beltsville, MD 20705-2351. Retain one copy for your files. All items on the face of the application are self explanatory unless noted below. Corrections on the application form and exhibits must be initialed and dated. DO NOT use masking materials to make corrections. If a certificate is allowed, you will be requested to send a check payable to "Treasurer of the United States" in the amount of \$300 for issuance of the Certificate.

Plant Variety Protection Office Telephone: (301) 504-5518

ITEM

- 16a. Give: (1) the genealogy, include public and commercial varieties, lines, or clones used, and the breeding method;
 - (2) the details of subsequent stages of selection and multiplication;
 - (3) evidence of uniformity and stability; and
 - (4) the type and frequency of variants during reproduction and multiplication and state how these variants may be identified.
- 16b. Give a summary of the variety's distinctness. Clearly state how this application variety may be distinguished from all other varieties in the same crop. If the new variety is most similar to one variety or a group of related varieties:
 - (1) identify these varieties and state all differences objectively;
 - (2) attach statistical data for characters expressed numerically and demonstrate that these are clear differences;
 - (3) submit, if helpful, seed and plant specimens or photographs (prints) of seed and plant comparisons which clearly indicate distinctness.
- 16c. Exhibit C forms are available from the PVPO for most crops; specify crop kind, Fill in Exhibit C (Objective Description of the Variety) form as completely as possible to describe your variety.
- 16d. Optional additional characteristics and/or photographs. Describe any additional characteristics that cannot be accurately conveyed in Exhibit C. Use comparative varieties as is necessary to reveal more accurately the characteristics that are difficult to describe, such as plant habit, plant color, disease resistance, etc.
- 16e. Section 52(4) of the Act requires applicants to furnish a statement of the basis of the applicant's ownership. The applicant may be the actual breeder, the employee of the breeder, the owner through purchase of inheritance, etc.
- 17. If "yes" is specified (seed of this variety be sold by a variety name only, as a class of certified seed), the applicant may NOT reverse this affirmative decision after the variety has been sold and so labelled, the decision published, or the certificate issued. However, if "No" has been specified, the applicant may change the choice. (See P.L. 103-349 for additional information.)
- 20. See Sections 41, 42, and 43 of the Act and Section 97.175 of the regulations for eligibility requirements.

NOTES: It is the responsibility of the applicant/owner to keep the PVPO informed of any changes of address or change of ownership or assignment during the life of the application/certificate. There is no charge for filing a change of address. The fee for filing a change of ownership or assignment is specified in Section 97.175 of the regulations. (See Section 101 of the Act, and Sections 97.130, 97.131, 97.175(h) of Regulations and Rules of Practice.)

To avoid conflict with other variety names in use, the applicant should check the variety names proposed by contacting: Seed Branch, AMS, USDA, Room 213, Building 306, Beltsville Agricultural Research Center—East, Beltsville, MD 20705. Telephone: (301) 504-8089.

Public reporting burden for this collection of information is estimated to average 30 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Department of Agriculture, Clearance Officer, OIRM, AG Box 7630, Washington, DC 20250; and to the Office of Management and Budget, Paperwork Reduction Project (OMB No. 0581-0055), Washington, DC 20503.

EXHIBIT A. ORIGIN AND BREEDING HISTORY OF THE VARIETY

VARIETY: TORONTO

Genealogy/Details of stages of selection and multiplication:

Toronto was developed from a cross between Salinas-88 (USDA, Salinas) and a breeding line, made in 1990. This line arose from a cross between an F5-line from which the variety Adal (Rijk Zwaan) was selected and another cross. This cross was made between Marbello (Bruinsma, the Netherlands) and a greenhouse butterhead lettuce breeding line of TS Seeds. A combined line and pedigree selection resulted in 1996 in a uniform F7-line based on one F6plant. All crossing and breeding was performed in the Netherlands. Toronto obtained the introduction number 45-18 RZ, which was changed to Toronto during the European registration procedure.

Selection was based on:

- 1. Field evaluation:
 - Homogeneity
 - Plant size
 - Head size
 - Head firmness
 - Head weight
 - Tipburn resistance
 - Bolting resistance
- 2. Laboratory tests on downy mildew and root aphid resistance, and LMV tolerance

Statement of uniformity and stability

Toronto was stable and uniform in variety tests and in seed multiplications carried out since 1996. Official tests of NAKTuinbouw (EC Variety List) and Plant Research International (Community Plant Variety Right) confirmed the uniformity and stability of Toronto.

Variants In seed multiplications we are finding about 0.4% of off-types. These off-types are are big wild plants that do not make a head.




Fig. 1 Pedigree of lettuce variety 'Toronto'

Exhibit A

The number of generations over which uniformity and stability has been observed is: 2. Uniformity and stability have been shown since 1996.

Toronto is most similar to Salinas 88.

root aphid resistance - see remark in e-mail of December 30, 2004 on leftuce appl. 200000214 'Adal'.

LMV (lettuce mosaic virus) - Toronto contains the recessive mo1.2-gene, which provides resistance to the common LMV-0-strain and the LMV-1-strain (Pink et al., 1992. Euphytica 63: 169-174). This gene was originally described by Ryder (Journal of the ASHS 95, 378-379. 1970). It was recently cloned by Nicaise et al. (Plant Physiol 132: 1272-1282). This gene is absent in the comparison varieties Fortunas, Vetonas, and Adal.

Trial 1

location: Hendrik-Ido-Ambacht, the Netherlands

sowing: 17-03-1998 planting: 27-04-1998 harvest: 22-06-1998

variety: Fiorette(n=10) Toronto(n=10) Adal(n=10) days to bolting: 163.4(sd=1.2) 153.0(sd=1.2) 163.0(sd=1.3)

variety: Fiorette (n=10) Toronto(n=10) color (RHS-color chart) 146C 146A

Trial 2

location: Roelofarendsveen, the Netherlands

sowing: 19-03-1999 planting: 20-04-1999 harvest: 16-06-1999

variety: Fiorette(n=10) Toronto(n=10) Adal(n=10) days to bolting: 150.7(sd=1.5) 140.7(sd=1.2) 151.1(sd=1.8)

variety: Fiorette (n=10) Toronto (n=10) color (RHS-color chart) 146C 146A

F-tests based on analysis of variance across both locations show that Toronto is significantly different from Fiorette for days to bolting (F=960; p<0.001). F-tests based on analysis of variance across both locations show that Toronto is significantly different from Adal for days to bolting (F=742; p<0.001).

As stated in the original exhibit B, 'Toronto' is different from 'Salinas-88' by its resistance against Bremia lactucae strains NL3, NL4, NL10, NL11, NL12, NL13, NL15, NL16, and Bl:21. This resistance is provided by theDm18-gene, but we are not sure that this is the only Bremia resistance gene in 'Toronto'.

All of the lettuce downy mildew (Bremia lactucae) Â strains NL3, NL4, NL10, NL11, NL12, NL13, NL15, NL16, and Bl:21 are controlled by the Dm18-gene.

The following can be added about the resistance of Toronto:
'Toronto' is most similar to 'Salinas 88'; however, 'Toronto' differs from 'Salinas 88' in resistance to lettuce downy mildew (Bremia lactucae) pathotypes CA I, CA 2a, CA 2b, CA III, CA IV, CAV, and CA VI and carries the DM18 gene whereas 'Salinas 88' does not contain the DM18 gene and is susceptible.

EXHIBIT B. STATEMENT OF DISTINCTNESS

VARIETY: TORONTO

Identify the varieties and state all differences objectively:

Toronto is a slow bolting crisp lettuce with LMV resistance. Toronto is resistant to the official Bremia races NL1, NL2, NL3, NL4, NL5, NL6, NL7, NL10, NL11, NL12, NL13, NL14, NL15, NL16, B1-19 and B1-21.

Difference from other lettuce varieties from the database:

| Denomination of parent variety | Characteristic in which the similar variety is different | State of expression of similar variety | State of expression of candidate variety |
|------------------------------------|--|---|---|
| Adal | root aphid resistance (Pemphigus bursarius) | resistant | susceptible |
| Salinas 88 Vetonas, Fortunas | downy mildew resistance (<i>Bremia lactucae</i>) | resistant to NL1, NL2, NL5, NL6, NL7, NL14, Bl-19 | resistant to NL1, NL2, NL3, NL4, NL5, NL6, NL7, NL10, NL11, NL12, NL13, NL14, NL15, NL16, B L19; Bl-21 |
| Adal | downy mildew resistance (<i>Bremia lactucae</i>) | resistant to NL1, NL2, NL4, NL5, NL6, NL13, NL14, NL15, BI-17, -BI-49- | resistant to NL1, NL2, NL3, NL4, NL5, NL6, NL7, NL10, NL11, NL12, NL13, NL14, NL15, NL16, Bl 19, Bl-21 |
| Vetonas, Fortunas | lettuce aphid resistance (Nasonovia ribisnigri) | resistant | susceptible |
| Adal, Fortunas, Vetonas | LMV resistance | susceptible | resistant |
| Fiorette, Adal | bolting resistance | slower bolting | faster bolting |
| Fiorette | leaf dentation | 2 shallowly dentate | 4 crenate |
| Fiorette | leaf colour intensity | 6 medium to dark | 7 dark |

4/10/06

U.S. Department of Agriculture Agricultural Marketing Service Livestock and Seed Division **OBJECTIVE DESCRIPTION OF VARIETY**

| | | | LETTUCE Lactuca s | ativa | | |
|-----------------------|---|--|---------------------------------------|------------------------------------|-----------------------|--|
| | APPLICANT(S) | | | FOR O | FFICIAL USE ONLY | |
| Rijk Zwaar | n Zaadteelt en Zaadhand | el B.V. | PVPO N | UMBER 20 | 000021 | 5 |
| ADDRESS | (Street and No. or R.F.D. No., C | City, State, and ZIP-Code) | . VARIET | Y NAME | | |
| Burgemees | ster Crezeelaan 40 | | TORON | ТО | | |
| 2678 KX The Nether | | | EXPERIM | MENTAL DESIGNATION | | |
| | | e characters which best o | | | | oriate number (at leas |
| | on of the test area is: | orabolitariai obolitty tir airy | | stem Used: | THE PLANE COLO. | |
| The Nether | | sted check varieties page | RHS | | | |
| 0 6 | 01 = Cutting/Leaf 02 = Butterhead 03 = Bibb | oted check varieties page 05 = Great Lakes Gr 06 = Vanguard Group 07 = Imperial Group | oup 09=Stem | | | |
| | 04 = Cos or Romaine | 08 = Eastern (Ithaca |) Group | | | nonana a assembla an amisino a amisino dia anto colora a colora de colora a colora a colora a colora a colora a |
| 2. SEED: | COLOR | LIGHT DORM | | AT DORMANCY | | |
| | 1 = White(Silver Gray) 2 = Black (Gray Brown | | | = Susceptible = Not susceptible | | |
| 2 | 3 = Brown(Amber) | | . | • | | |
| | | | | | | |
| 3. COTYLE | DON TO FOURTH LEAF | STAGE: NOTE: Provide a | | tocopy of the fourth lea | f from 20 day old see | dling |
| | | grown un | der optimal conditions | | 4 | |
| 3 | SHAPE OF COTYLED | ONS: 1=Broad 2=In | termediate 3 = Spatul | ate | | |
| | | ···· | | | | |
| 3 | SHAPE OF FOURTH | LEAF: | | | | |
| | | \wedge | Λ | | | • |
| | | | /// 조1건 | | | |
| | | (| 111 513 | | | |
| | | | (1) 3/5 | | | |
| | 1 2 | 3 4 | (I) 45 | | | |
| 1 7 |] | | 5 | | | |
| 1 / | LENGTH/WIDTH INDI | EX OF FOURTH LEAF: L/V | V x 10 | | | |
| 2 | · . | | | | | <u>and a state of the first of th</u> |
| 3 | APICAL MARGIN: | 1 = Entire | 4 = Moderately Dense | 7 = Lobed | | |
| | DACAL MARDOIN | 2 = Creanate/Gnawed | 5 = Coarsely Dentate | 8 = OTHER (specify) | | |
| 3 | BASAL MARGIN: | 3 = Finely Dentate | 6 = Incised | | | |
| | LINEDLY ATION | | | | | |
| 2 | UNDULATION: | 1 = Flat | 2 = Slight | 3 ≔ Medium | 4 = Marked | |
| · <u> </u> | | | · · · · · · · · · · · · · · · · · · · | | | <u> </u> |
| 4 | GREEN COLOR: | 1 = Yellow Green | 3=Medium Green | 5 = Blue Green | 7 = Gray Green | |
| | GREEN COLON. | ,2 = Light Green | 4 = Dark Green | 6 = Silver Green | 7 - dray dreen | ** * |
| 4 | · | | | | | <u> </u> |
| | ANTHOCYANIN: | | | | • | e e |
| 1 | | | | | | |
| | DISTRIBUTION: | 1 = Absent 2 = Margin Only | 3 = Spotted 4 = Throughout | 5 = OTHER (specify) | | * |
| | , | - margar only | - Throughout | | | |
| | CONCENTRATION: | 1 = Light | 2 = Moderate | 3 = Intense | | |
| · | | | | | | |
| 2 | ROLLING: | 1 = Absent | 2 = Present | | | |
| | | | | | | |
| 1 | CUPPING: | 1 = Uncupped | 2 = Slight | 3 = Markedly | | |
| | | | · | | | <u> </u> |
| 3 | REFLEXING: | 1 = None | 2 = Apical margin | 3 = Lateral Margin | | |
| | | | | | | |

| - | L | (finest divisions of the margin | | | | | 3 = 6+00 | ong (Great Lakes 65 | | | |
|-----------|------------|--|---|---|--|---------------------|---------------------|---------------------------------------|--|--|--|
| | 2 | UNDULATION OF THE APICAL MARGIN: | 1 = Absent/Slight | (Dark Green Bostor | a) 2 = Moderate (Vanguard) | | 3 = Stro | ong (Great Lakes 65 | | | |
| | 4 | GREEN COLOR: | 1=Very Light Gre 2=Light Green (N | | 3 = Medium Green (Great 4 = Dark Green (Vanguard | | | | | | |
| | 1 | ANTHOCYANIN (grown DISTRIBUTION: | at or below 10 °C): 1 = Absent 2 = Margin Only (I | Big Boston) | 3 = Spotted (Calif. Cream 4 = Throughout (Prize Hea | | 5 = OTHER (specify) | | | | |
| | | CONCENTRATION: | 1 = Light (Iceberg) | | 2 = Moderate (Prize Head) | | 3 = Inte | ense (Ruby) | | | |
| | 3 | SIZE: | 1 = Small | | 2 = Medium | *** | 3 = Larg | ge | | | |
| | 1 | GLOSSINESS: | 1 = Dull (Vanguard |) | 2 = Moderate (Salinas) | | 3 = Glos | ssy (Great Lakes) | | | |
| | 1 | BLISTERING: | 1 = Absent/Slight (| Salinas) | 2 = Moderate (Vanguard) | | 3 = Stro | ong (Prize Head) | | | |
| | 3 | LEAF THICKNESS: | 1 = Thin | | 2 = Intermediate | | 3=Thi | ck | | | |
| | 1 | TRICHOMES: | 1 = Absent (smooth | ·) | 2 = Present (spiny) | 2 = Present (spiny) | | | | | |
| . PLAN | IT (at ma | rket stage. Choose a comparison va | | type): | | , | | | | | |
| 4 | 0 | SPREAD OF FRAME LE. | AVES: | 3 8 | cm. Salinas | (spec | ify compai | rison variety) | | | |
| 1 | 5 | HEAD DIAMETER (market cm. This variety | trimmed with single cap | leaf): 1 5 | cm. <u>Salinas</u> | (speci | fy compar | ison variety) | | | |
| | 2 | 1 | 1 = Flattened 2 = Slightly Flatten | ed | 3 = Spherical 5 = Elongate | | | on-heading OTHER | | | |
| . [| 3 | HEAD SIZE CLASS: | 1 = Small | | 2 = Medium | | 3=L | arge | | | |
| | | HEAD COUNT PER CARTO | ON | | | | | · | | | |
| | 8 | 5 4 HEAD W g This Va | - | 8 6 5 | g. Salinas (specify con | nparison variety) | | | | | |
| | 4 | · · | I = Loose 2 = Moderate | | 3=Firm 4=Very Firm | | | | | | |
| BUTT F | , | of market-trimmed head): | | | | | | | | | |
| | 3 | SHAPE: 1 | = Slightly Concav | е | 2 = Flat | unded | | | | | |
| | 2 | MIDRIB: 5 1 | = Flattened (Salina | es) | 2 = Moderately Raised | | | minently Raised eat Lakes 659) | | | |
| | · · · · · | market-trimmed head): mm Diameter at base | of Head | | | | | | | | |
| 3 | 4 | 7 Ratio of head | diameter/core diam | eter | | | | | | | |
| 4 | 2 | Core height from base of mm. This Variety | of head to apex: | 3 9 m | ım. <u>Salinas</u> (specify co | mparison variety) | | (| | | |
| BOLTI | ING (Gi | | 19/1999): Note: Firs | | eed first receives adequate moisture | | | : | | | |
| | 4 | | to germir | ate. This can and offen d | desi'equal the planting date. nergence (summer conditions): | 4 | 7 | specify comparison variety Salinas | | | |
| 1 | 4 | TITUS VAILELY | | | | 1 | 5 = Very Rapid | | | | |
| . 1 | <u>" </u> | 1 BOLTING CLASS | S:1=Very Slow 2= Slow | 13.00 (1.1.1.1.1.1.1.1.1.1.1.1.1.1.1.1.1.1.1. | 3—Medium 4≐ Rapid | • | 5 = Ver | y Rapid | | | |

| | | | | | | | | | | | | | | | | | 20 | 0 (| 0 (| 0 2 | 9 | 5 | |
|----------|----------|------------------------|---------------------|-----------------|----------|---------|----------|-----------|------|--|-----------|------------|----------|--------------------|---------------|----------|---------|-------------|---------|--------------------|-------|--|---|
| 3 | 0 | | ad of B This Va | | ant | (at wid | est poir | nt): | | 3 | 0 | cm. | Salir | nas | | (specify | | on variety | | | , 8 | | |
| | 2 | BOLT | ER LEA | VES: | | , | 1 = .5 | Straigh | t | | | , | 2 | = Cı | urved | | | | | • | | | |
| | 2 | | /ARGIN | : | | | 1 = E | ntire | | | | | 2 | = D | entate | | | | | | | | |
| | 2 | | OLOR: | | | | 1 = I | ight G | irec | en ' | , | | 2 | = N | Medium Gre | en | | 3 = | = Dark | Green | 1 | | |
| | L | ــ | R HAB | T: | | | | | | | | | | | | | | | | | | | |
| | 2 | , TE | RMINA FLORES | L | : | | 1 = | Absen | t | | | | 2 | = Pr | resent | | _ | | | | | <u>. </u> | |
| | 2 | 1 | ΓERAL (re head) | SHOOT | S | | 1 = 7 | Absent | : | | | | 2 | = P | resent | | | | | | | | |
| | 2 | BA | SAL SIE | E SHO | отя | s | 1 = A | Absent | - | | | | 2 | = Pr | resent | | | | | | | | |
| | | Y <i>(earliness d</i> | | | | | | eason. | | | | | | | | | | | | | | | |
| | ASON | Applic. 1 | | | | | | days | | | | | | | CHEC | Κ VΔΕ | RIETY 2 | 9/ | <u></u> | | | | |
| J. | 13011 | Аррис. | 7# Ol u | <u>ays</u>] | Ĭ | IIGUK | 1,7 0 | uays | ı | | | - | | | CHEC | K VAI | 11611 4 | -1 | | | | | |
| Spr | ing | | | | | | | | | | | | | | | | | | | | | | |
| | | 9 | 4 | | | | 9 | 6 | | | | | | | | | * | | | | | | |
| Sun | nmer | | | - | | | | | | Salina | ıs | | | | | • | | | | | * | - | _ |
| Fall | | | | | | | | | | | | | | | | | | | | | | | |
| | | <u> </u> | | 1 | | | | | | | | | | | | | | | | | | | _ |
| Win | iter | | | | Ц | | | | | <u>. </u> | | | | | | | | | | | | i | |
| Give p | lanting | date(s), a | nd loca | tion(s): | ٠ | | | | | | | | | | | | | | | | | | |
| Spring | | | • | | | | | | | | | | | | | | | | | | | | |
| Summe | er C | 3/19/199 | 9. Roel | ofarend | sve | en tl | ne Ne | therlan | ıds | | | | | • | • | | | | | | | | |
| Fall | | .0,.0,.00 | | 01010 | | , | 10 110 | | - | | | | | | | | | | | | | | _ |
| | | | | | | | | | | | | | | | | | ** | | | | | | _ |
| Winter | | | | | | | | | | | | | | | : | | | | | | | | _ |
| 1/ First | water | date to h | arvest. | | 2 | 2/ Fill | in ch | eck va | riet | y nam | e on ti | he app | ropriate | e lin o |). | | | | | | | | |
| 10. AD | APTA | ΓΙΟΝ: | | | | | | | | | | | • | | | | | | | | | | |
| | | PRIMAR | Y REGIO | ONS OF | ΑE | DAPTA | ATION | l (tested | l an | d proven | adapted | <i>():</i> | (| 0 = | Not tested | | 1 = N | ot Adap | ted | 2 | = Ada | ipted) | |
| | 0 | South | west (C | alif., Ar | iz. | deser | t) . | 0 | | West | Coast | | 0 | | Northeast | | | | | | | | |
| | 0 | North | entral | '; | <i>,</i> | | | 0 | | South | ıeast | | | 7 | OTHER | | | | | | | | |
| | <u> </u> | SEASO | | | | Casis | • | | | | 1 | | | | Eall (area | Cnain | | | |) | | | |
| | - | 2 | Shiii | ng (area | | Span | 1 | | | | ' | | 2 | | Fall (area | эрап | | | | , | | | |
| | | | Sum | mer (ar | ea_ | Spai | n | | | | _) | | | | Winter (ar | ea Sp | ain | | |) | | | |
| | | | | | _ | | | | | | | | 1 | | | | | | | | | | |
| | 1 | GREEN | HOUSE | : | ı | 0 = 1 | Vot te | sted | | | | | | 1 | = Not Ada | pted | | <u> </u> | 2 | = Ada _l | pted | | |
| | | SOIL TYPE: 1 = Mineral | | | | | | | | 2 : | = Organic | | | | 3 : | = Both | | | _ | | | | |


| 11. DISEASE | S AND S | TRESS REACTIONS (0 = Not tested; 1 = | = Susceptible; | 2 = Intermediate; 3 = | = Resista | 2000 int; 4 = Highly | UUZ resistant; 5 | Tolerant): |
|-------------|---------|--------------------------------------|----------------|-----------------------|-----------|-------------------------|---------------------|------------------|
| | | <u>VIRUS</u> | | FUN | IGAL/BA | CTERIAL | | |
| | 0 | Big Vein | 1 | Corky Root Rot (Pyt | thium Ro | ot Rot) | | |
| | 3 | Lettuce Mosaic | 3 | Downy Mildew (Rad | ces see E | xhibit B) | | _ |
| | 0 | Cucumber Mosaic | 0 | Powdery Mildew | | | | |
| | 0 | Broad Bean Wilt | . 0 | Sclerotinia Rot | | | | |
| | 0 | Turnip Mosaic | 0 | Bacterial Soft Rot (F | Pseudome | onas spp.&oth | ers) | |
| | Ò. | Beet Western Yellows | 0 | Botrytis (Gray Mold) |) | | | |
| | 0 | Lett. Infectious Yellows | 0 | OTHER | | | <u> </u> | <u> </u> |
| | 0 | Other Virus | | | | , | | |
| | | I <u>NSECTS</u> | | <u>РН)</u> | YSIOLOG | ÎĈAL/STRESS | | |
| | 0 | Cabbage Loopers | 0 | Tip Burn | 0 | Salt | | |
| | 1 | Root Aphids | 2 | Heat | 2 | Brown Rib (F | Rib Discolorat | tion, Rib Blight |
| | 0 | Green Peach Aphid | 0 | Drought | | OTHER | | |
| | 0 | Other Insect | 0 | Cold | |] 0141611 | | <u> </u> |
| | | | | 5514 | | | | • |
| | | POST HARVEST | | | | | ŧ | · · |
| | 0 | Pink Rib | 0 | Internal Rib Necros | is (Black | heart, Gray Rib | o, Gray Streat | k) |
| | 0 | Russet Spotting | 0 | Brown Stain | | | | |
| ř. | 0 | Rusty Brown Discoloration | | | | | | |
| 12. BIOCHEM | ICAL OR | ELECTROPHORETIC MARKERS: | | | | | | |
| 13. COMMEN | TS: | | | | | | | |
| | | | | | | | | |
| | | · · | | | | | | |
| | | suc | GGESTED CHE | CK VARIETIES | | | | |
| • | | | | | | | | |



45-18 RZ Toronto' Mature leaf



45-18 RZ Toronto' Mature leaf, from above



45-18 RZ Toronto' 4th leaf

EXHIBIT D. ADDITIONAL DESCRIPTION OF THE VARIETY

VARIETY: TORONTO

Toronto is a crisp lettuce with LMV resistance. It makes medium sized heads with a very stable round shape. Toronto is slow bolting. Toronto is resistant to the official Bremia races NL1, NL2, NL3, NL4, NL5, NL6, NL7, NL10, NL11, NL12, NL13, NL14, NL15, NL16, Bl-15 and Bl-21 and the Californian pathotypes Ca I, Ca-IIA, Ca-IIB, Ca-IV, and Ca V.

RAD 060

Toronto can be used in plantings from Spring until Fall

| REPRODUCE LOCALLY. Include form number and date on all reproductions. U.S. DEPARTMENT OF AGRICULTURE | FORM APPROVED - OMB NO. 0581-0055 EXPIRES: 12-31-9 |
|--|---|
| AGRICULTURAL MARKETING SERVICE | The following statements are made in accordance with the Privacy Act of 1974 (5 U.S.C. 552a) and the Paperwork Reduction Act (PRA) of 1995. |
| SCIENCE AND TECHNOLOGY DIVISION - PLANT VARIETY PROTECTION OFFICE | Application is required in order to determine if a plant variety protection |
| EXHIBIT E STATEMENT OF THE BASIS OF OWNERSHIP | certificate is to be issued (7 U.S.C. 2421). Information is held confidentia until certificate is issued (7 U.S.C. 2426). |
| 1. NAME OF APPLICANT(S) | |
| Rijk Zwaan Zaadteelt en Zaadhandel B.V. | 2. TEMPORARY DESIGNATION 3. VARIETY NAME OR EXPERIMENTAL NUMBER |
| Kijk Swaan Saadteelt en Saadnandel B.v. | |
| | TORONTO |
| 4. ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP Code, and Country) | 5. TELEPHONE (include area code) 6. FAX (include area code) |
| Burgemeester Crezeelaan 40 | + 31 174 532300 + 31 174 513730 |
| 2678 KX DE LIER, The Netherlands | 7. PVPO NUMBER |
| | 200000215 |
| 8. Does the applicant own all rights to the variety? Mark an "X" in appropriate | e block. If no, please explain. x YES NO |
| | |
| | |
| | |
| | |
| | |
| 9. Is the applicant (individual or company) a U.S. national or U.S. based compa | any? YES NO RAD |
| If no, give name of country <u>Netherlands</u> | 4/10/8 |
| 10. Is the applicant the original breeder? If no, please answer the following: | |
| Waling the American control of the C | X YES NO |
| a. If original rights to variety were owned by individual(s): ls (are) the original breeder(s) a U.S. national(s)? If no, give name o | of country |
| | |
| | |
| | YES NO |
| b. If original rights to variety were owned by a company: | |
| is the original breeder(s) U.S. based company? If no, give name of o | country |
| | |
| 11. Additional explantion on ownership (If needed, use reverse for extra space) | l: |
| | |
| | |
| | |
| LEASE NOTE: | |
| lant variety protection can be afforded only to owners (not licensees) who meet | t one of the following criteria: |
| sandly protoction out to antorada only to aminota that meanaces? Wite mean | . One of the following effection |
| . If the rights to the variety are owned by the original breeder, that person must of a country which affords similar protection to nationals of the U.S. for the s | · · · · · · · · · · · · · · · · · · · |
| If the rights to the variety are owned by the company which employed the or nationals of a UPOV member country, or owned by nationals of a country whi genus and species. | |
| . If the applicant is an owner who is not the original breeder, both the original b | preeder and the applicant must meet one of the above criteria. |
| ne original breeder may be the individual or company who directed final breefinition. | eding. See Section 41(a)(2) of the Plant Variety Protection Act for |

Public reporting burden for this collection of information is estimated to average 10 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Department of Agriculture, Clearance Officer, OIRM, AG Box 7630, Jamie L. Whitten Building, Washington, D.C. 20250. When replying, refer to OMB No. 0581-0055 and form number in your letter.

Under the PRA of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

The U.S. Department of Agriculture (USDA) prohibits discrimination in its programs on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, and marital or familial status. (Not all prohibited bases apply to all programs). Persons with disabilities who require alternative means for communication of program information (braille, large print, audiotape, etc.) should contact the USDA Office of Communications at (202) 720-2791.

To file a complaint, write the Secretary of Agriculture, U.S. Department of Agriculture, Washington, D.C. 20250, or call (202) 720-7327 (voice) or (202) 720-1127 (TDD). USDA is an equal employment opportunity employer.